

Supplemental Table 6: Autism spectrum disorder symptoms

Assessment		Classification		
Domain	Description; range; M (SD)	Met criteria; n (%)	ASD Diagnosis/Met classification; n(%)	
ASD Consensus				
	Based on ADOS-2, ADI and clinical evaluation (DSM-5)		5 (24)	
Psychiatric Evaluation				
<i>Social communication and interaction deficits (n=21)</i>	Deficits in socio-emotional reciprocity	8 (38)	5 (35)	5(24)
	Deficits in non-verbal communication	5 (24)		
	Deficits in developing and maintaining friendships	20 (95)		
<i>Repetitive Behaviors (n=11)</i>	Stereotyped behaviors	17 (80)	20(95)	
	Insistence of sameness	15 (71)		
	Highly restricted interests	16 (76)		
	Sensory symptoms	20 (95)		
ADOS-2				
<i>Social affect calibrated scores (n=19)</i>	1-3	5.1(1.77)	5 (26)	12(57)
	4-5		7 (37)	
	6-10		7 (37)	
<i>Repetitive behavior calibrated score (n=19)</i>	1-3	6.33(2.52)	2 (11)	
	4-5		4 (21)	
	6-10		13 (68)	
<i>ADOS comparison scores (n=21)</i>	No ASD (1-3)	5.25 (1.94)	9 (43)	
	ASD (4-5)		2 (10)	
	Autism (6-10)		10 (47)	
ADI-R				
<i>Socialization (n=21)</i>	3 – 26	13.1 (7.06)	13 (61)	10(47)
<i>Communication (n=21)</i>	2 – 21	10.6 (5.04)	16 (76)	
<i>Repetitive Behaviors (n=21)</i>	0 – 9	4.6 (2.87)	15 (71)	